Presenters and Poster Titles

Emporia State University

Ryan Frier	WAVELET-BASED ACOUSTIC CLASSIFICATION OF BIRD SPECIES
Thomas Houk	IMPROVED TRACKING OF KANSAS ECONOMIC ACTIVITY
Mary M. Ralston	DNA BARCODING OF BLOW FLY SPECIES IN COFFEY COUNTY, KANSAS
Janelle Thimmesch	IDENTIFICATION AND QUANTIFICATION OF IBUPROFEN IN BLOOD USNG HPLC-UV/VIS
Randol Wehrbein	FOSSIL OF THE OLDEST HEALING BONE FRACTURE IN AMNIOTES

Rachel Bechtold	BACTERIAL DIVERSITY OF AN ABANDONED MINE LAND SOIL IN SOUTHEAST KANSAS
Sanket Bhoyate	GREEN POLYURETHANES FOR INDUSTRIAL APPLICATIONS USING ECO- RENEWABLE RESOURCES
Lynzee Flores	EMOTION ETQq0030 0 396 612 reW*n51 gR0.051 g/ 612 re.

Heather V. Nelson-Brantley	IMPROVING RURAL CRITICAL ACCESS HOSPITAL OUTCOMES: AN EXEMPLARY CASE STUDY
Joshua Breeden	INHIBITION OF STAT3 SUPPRESSES ANGIOTENSIN II-INDUCED CARDIOMYOBLAST HYPERTROPHY THROUGH MODULATING THE PATHWAY
Ian Huck	IDENTIFICATION OF HEPATOCYTE NUCLEAR FACTOR 4 ALPHA (HNF4) TARGET GENE SIGNATURE AND VALIDATION AS A PROGNOSTIC BIOMARKER OF HEPATOCELLULAR CARCINOMA
Jacob New	SECRETORY AUTOPHAGY IN TUMOR ASSOCIATED FIBROBLASTS PROMOTES HEAD AND NECK SQUAMOUS CELL CARCINOMA PROGRESSION AND EMERGES AS A NOVEL THERAPEUTIC TARGET
Kelly Zarifa	TO HAVE A VOICE: TREATMENT PLANNING FOR PERSONS WITH APHASIA, THE KANSAS CONNECTION

Wichita State University

Mahmood Al Bashir	FEMTOSECOND LASER NANOTEXTURING OF DRUG-ELUTING STENTS
Daniel Clifford	REDUCING PTSD FOR VETERANS AND EMERGENCY SERVICE PROVIDERS
Kelsay E. Gardiepy	IT BLEEDS, IT LEADS #HESSTONSTRONG: A FANTASY-THEME ANALYSIS OF LOCAL AND NATIONAL NEWS COVERAGE OF THE 2016 HESSTON, KANSAS SHOOTING
Alicia Keow	SOLAR ENERGY HARVESTING USING IONIC POLYMER-METAL ENHANCED WATER ELECTROLYSIS
Mostafa Hasan	DETERMINING AN EFFECTIVE TREATMENT PLAN FOR BREAST CANCER: A MULTI-CRITERIA DECISION MODEL AND ALGORITHM
Lipilekha Mukherjee	DIFFERENTIAL REGULATORY GENE EXPRESSION AT THE PROTEOMIC LEVEL IN SUBTYPES OF HUMAN OVARIAN CANCER CELL-LINES
Erin Sawyers	NUTRITION GUIDELINES IN KANSAS ELEMENTARY SCHOOLS
Akshay Reddy Tummala	CUBESAT PLATFORM FOR RESEARCH, EDUCATION AND BUSINESS IN KANSAS
Mercy Umeri	IMPROVING TRANSIT IN WICHITA: MARRYING THE OLD AND THE NEW

WAVELET-BASED ACOUSTIC CLASSIFICATION OF BIRD SPECIES

Ryan Frier and Qiang Shi
Department of Mathematics and Economics, Emporia State University

In this work, we use wavelet packet decomposition to classify bird species based off their calls. Knowing the bird species present in a given habitat can tell us many important things about the area. For instance, knowing the bird species helps us track the extinction of birds. Also, since birds are sensitive to changes in the ecosystem, knowing how many bird species are present in an ecosystem tells us how healthy the system is. The current methods of determining bird species have their limitations. Those methods usually need highly trained ornithologists to identify size, shape, posture, field marks, habitat, and flight patterns. The process requires many hours, and the results may be subjective. In our study, we developed a semi-automatic classification algorithm by using wavelet transformations, a mathematical tool which has been successfully applied in many areas. For example, the FBI uses a wavelet-based scheme to compress their fingerprint images. We consider the calls of the Whip-poor-will, the Northern Bobwhite, the Barred Owl, the Eastern Kingbird, and the Common Raven. We first segment the bird calls into syllables. Then we apply wavelet decomposition to decompose those syllables and extract certain parameters from the transformed signals. All of the instances and the parameters are compiled into a training set and a test set. We then classify the instances. Among all the classifiers we tested, Random Tree and Random Forest were the most successful. Both classifiers achieved over 70% accuracy.

IMPROVED TRACKING OF KANSAS ECONOMIC ACTIVITY

Thomas Houk and Marc Fusaro
School of Business, Emporia State University

Forecasting state level gross domestic product (GDP) has been notoriously difficult as past Kansas revenue over-projections have highlighted. This research develops a better estimating method for state GDP growth rates, at higher frequency, and more timely than current data provided by the Bureau of Economic Analysis (BEA). While the data provided by the U.S. Department of Commerce is reliable, the release cycles for quarterly state GDP can be as lengthy as five months before being reported. This research provides estimates on state level GDP growth at monthly frequencies with release cycles as early as five weeks after the end of each month. The methodology relies on time series forecasting on data collected from the Bureau of Labor Statistics, BEA, and Zillow Research. An ARIMA regression model is applied on historical data as well as the last twenty-four periods to generate an estimate on growth for recent months. The research borrows

DNA BARCODING OF BLOW FLY SPECIES IN COFFEY COUNTY, KANSAS

Mary M. Ralston and Scott S. Crupper
Department of Biological Sciences, Emporia State University

Blow flies are one of the first insects to arrive during decomposition, thereby allowing forensic

FOSSIL OF THE OLDEST HEALING BONE FRACTURE IN AMNIOTES

Randol Wehrbein[1], Michael Morales[1], and A.P. McElroy [2]

[1]Department of Physical Sciences, Emporia State University; [2]Kansas Department of Health and Environment

In Greenwood County, near Toronto, Kansas, a fossil locality in the Snyderville (Shale) Member of the Oread (Limestone) Formation (late Pennsylvanian Period, c. 300 million years ago) contains many disarticulated and broken vertebrate fossilized bones weathering out of an ancient soil. A notable specimen is a bone from an early amniote that was fractured while the animal was alive and then partially healed prior to its death. Amniotes are vertebrate animals that lay shelled eggs

THE RELATIONSHIP BETWEEN PSYCHOPATHY AND PERCEPTIONS OF OTHERS

Arianne Fisher and W. Trey Hill
Department of Psychology, Fort Hays State University

Psychopathy is a complex construct surrounded by many unanswered questions. It is characterized by dominance, superficial charm, manipulation, impulsivity, and a lack of empathy and remorse (Hare & Neumann, 2009). Additionally, psychopathy is often associated with many antisocial and criminal behaviors. This is thought to be the result of the characteristic lack of empathy. Therefore, it seems prudent to attempt to increase empathy levels in this population in order to decrease these often-harmful behaviors. One empathy-inducing literature intervention has shown promise in general populations in the past (Johnson, Cushman, Borden, & McCune, 2013). The present study attempted to apply this intervention to increase empathy levels in those high in psychopathy. Ninety-
induce empathy and were given a psychopathy scale, an empathy scale, and a scale measuring perceptions of the characters. Half of the participants were taught to generate imagery while

Fort Hays

PHYLOGENETIC DISTRIBUTION OF AN ENDOGENOUS STRAIN OF DAHLIA MOSAIC VIRUS IN MEMBERS OF ASTERACEAE

Keri Caudle and Eric Gillock
Departments of Biological Sciences and Geosciences, Fort Hays State University

Dahlia Mosaic Virus (DMV) is a double-stranded DNA viral pathogen. A newly discovered strain of this virus called DMV-D10 was first observed in Dahlia variabilis. DMV-D10 does not induce visible symptoms of infection in the host plant, and is classified as an endogenous pararetrovirus. These viruses have the ability to integrate their viral sequences into the host plant genome, which can be transmitted vertically to offspring. Currently, no studies have examined the host range of DMV-D10. Because DMV-D10 has only been observed in Dahlia, the objective for this study was to determine if presence of DMV-D10 follows an evolutionary relationship among species closely related to Dahlia. It was hypothesized species in the same tribe (Coreopsideae) as Dahlia, such as Cosmos, were more likely to be infected with DMV-D10 compared to species within other tribes in the Asteraceae family. PCR results for a movement protein gene indicate DMV-D10 is more

EXPERIMENTAL NATURAL SELECTION OF BIG BLUESTEM ECOTYPES ACROSS THE GREAT PLAINS: A NOVEL TEST FOR THE STRENGTH OF LOCAL ADAPTATION

Matthew Galliart[1], Paul St Amand[2], Jesse Poland[3], Nora M. Bello[4], Mary Knapp[5], Sara G. Baer[6], David Gibson[6], Brian R. Maricle[7], and Loretta Johnson[1]

[1]Division of Biology, Kansas State University; [2]USDA, Hard Winter Wheat Genetics Research Unit, Kansas State University; [3]Department of Plant Pathology, Kansas State University; [4]Department of Statistics, Kansas State University; [5]Department of Agronomy, Kansas State University; [6]Plant Biology and Center for Ecology, Southern Illinois University; [7]

Kansas State University

BRINGING COMPUTATIONAL THINKING TO K-12
Joshua Levi Weese

EMOTION LABOR, PATIENT-CENTERED CARE & FIRST RESPONDERS DISASTER RESPONSE: LESSONS LEARNED FROM THE 2011 JOPLIN TORNADO

Flores, L., Ishmael H., Tims, M., Mason, A. & Drew, S.
Department of Communication, Pittsburg State University

The analysis examines the opportunities and obstacles experienced by first responders to providing patient-centered care within the first 48-hours following the EF-5 Joplin-Duquesne tornado that struck on May 22, 2011 in Joplin, Missouri. A total of 14 semi-structured qualitative interviews

Pittsburg

GLOBAL TRENDS IN SNOW COVER FREQUENCY

James Coll and Xingong Li
[1]*Department of Geography and Atmospheric Science, University of Kansas*

Trends in snow cover have yet to be qualitatively described in any meaningful way, particularly at the global level; despite the fact that snow melt can be a significant source of ground and surface water recharge. Additionally, estimates of the remaining lifespan of the high plains aquafer may fail to take into account the variable nature of climate and short-term trends in snow cover and precipitation, instead relying on background recharge rates which may no longer mirror surface water inputs. In this work, we present tshe magnitude and significance of snow cover frequency trends across the globe using more than 16 years of an improved and gap filled MODIS daily snow cover dataset from 10/01/2000 through 09/30/2016. This trend, available for exploration at trendy-snow.appspot.com, depicts a globally consistent and locally relevant snow cover assessment which allows Kansans and others to more accurately predict recharge rates; improve predictions of streamflow and water availability; validate local-, meso-, and macro-scale climatological forecasts; as a tool for winter planning and budgeting for municipalities; and to inform the general public about this critical aspect of the hydrologic cycle.

MALTREATMENT AND PHYSICAL HEALTH OUTCOMES IN FOSTER YOUTH: AN EXAMINATION OF PROTECTIVE FACTORS

Lindsay Huffhines
Clinical Child Psychology Program, University of Kansas

Child maltreatment and foster care placement are associated with physical health problems and high rates of health service use across the lifespan. Yet some individuals exposed to maltreatment and/or placed in foster care in childhood remain healthy. Protective factors, such as coping style, emotional/behavioral functioning, and foster family environment may explain differences in physical health outcomes. Therefore, the purpose of this research was to first describe the prevalence of chronic health conditions in a sample of 213 maltreated youth in foster care, enrolled

IDENTIFYING AREAS AT RISK FOR INJECTION-INDUCED SEISMICITY THROUGH SUBSURFACE ANALYSIS OF SOUTHERN KANSAS

Jeffrey C. Jennings

Department of Geology, University of Kansas

Kansas has experienced an unprecedented increase in seismicity linked to large-scale injection of wastewater. Subsurface faults and stresses, however, are not well known, making it difficult to properly site new disposal wells or make decisions regarding operations of existing wells. This study focuses on using publicly available well data to map subsurface faults, lineaments, and present-day stresses in southern Kansas, with the ultimate goal of identifying specific structures and regions that may be at higher risk for injection-induced seismicity. A database of >500,000 well tops from the Kansas Geological Survey was used to create new structure contour maps of 18 stratigraphic boundaries between the Precambrian basement rock and present-day surface.

INVESTIGATING THE IMPACT OF THE MUTATION CLUSTER REGION OF TUMOR-SUPPRESSOR ADENOMATOUS POLYPOSIS COLI (APC) ON COLON CARCINOGENESIS

Aaron J. Rudeen, Audrey L. Lamb, and Kristi L. Neufeld
Department of Molecular Biosciences, University of Kansas

Colorectal cancer is the third most commonly diagnosed cancer in both men and women. In Kansas, over 1,100 people will be diagnosed with colorectal cancer in 2016 alone. A vast majority (>80%) of colorectal cancers result from acquired genetic mutations to the tumor-suppressing protein Adenomatous Polyposis Coli (APC). Mutation of APC is widely considered to be an initiating event of colon carcinogenesis. Expression of mutant APC protein by cells leads to the deregulation of many cellular pathways and processes. The rapidly dividing cells of the small intestine and colon are especially susceptible to this disruption of normal cellular homeostasis, and eventually develop polyps. If not caught early or left untreated, these polyps give rise to aggressive carcinomas.

occur at this specific locus is not well understood, and is debated.

IMPROVING RURAL CRITICAL ACCESS HOSPITAL OUTCOMES: AN EXEMPLARY CASE STUDY

Heather V. Nelson-Brantley[1], Debra J. Ford[1], Karen L. Miller[1], Kristin Stegenga[1], Robert H. Lee[2], and Marjorie J. Bott[1]

[1]School of Nursing, University of Kansas Medical Center; [2]Department of Health Policy and Management, University of KaSchool o (,)6 (,)4 ()2 (Un)4aJ-140.21C5.83 -7.32Tc

IDENTIFICATION OF HEPATOCYTE NUCLEAR FACTOR 4 ALPHA (HNF4) TARGET GENE SIGNATURE AND VALIDATION AS A PROGNOSTIC BIOMARKER OF HEPATOCELLULAR CARCINOMA

Ian Huck[1], Chad Walesky[1], Sumedha Gunewardena[2], Steven Weinman[3], Timothy M. Schmitt[4], Sean C. Kumer[4], Udayan Apte[1]

[1]Department of Pharmacology, Toxicology and Therapeutics; [2]Department of Molecular and Integrative Physiology; [3]Department of Medicine; [4]Department of Surgery, University of Kansas Medical Center

Hepatocellular carcinoma (HCC) is the second leading cause of cancer-related death and the incidence of HCC is increasing in the US. HCC originates from hepatocytes, a cell responsible for many important liver functions such as carbohydrate and lipid metabolism, albumin and coagulation factor synthesis, bile acid synthesis, and xenobiotic metabolism. Hepatocytes are quiescent during normal conditions, but they possess enormous potential to proliferate and expand following liver injury. This proliferative capacity is beneficial during liver regeneration, but the mechanisms driving this cell division can contribute to HCC if left unchecked. Our lab studies a transcription factor known as Hepatocyte Nuclear Factor 4 alpha (HNF4) which is expressed at high levels in normal hepatocytes and inhibits hepatocyte proliferation. Our studies show that deletion of HNF4 in mice results in spontaneous proliferation and hepatic dedifferentiation. Further studied revealed that HNF4 deletion promotes HCC development in rodents and HNF4 expression is decreased in human HCC samples. Currently, early detection tools and accurate prognostic biomarkers for HCC are lacking. We identified an HNF4 target gene signature that can measure HNF4 activity. Using in silico data and in vivo mouse and human samples we demonstrate this signature is capable of distinguishing between early and late stages of HCC. In summary, our studies show that HNF4 is a therapeutic target in both regenerative medicine and cancer therapy and can be used as a prognostic biomarker in HCC. Our studies not only highlight an important biological mechanism but will have significant impact on treatment of liver cancers.

SECRETORY AUTOPHAGY IN TUMOR ASSOCIATED FIBROBLASTS PROMOTES HEAD AND NECK SQUAMOUS CELL CARCINOMA PROGRESSION AND EMERGES AS A NOVEL THERAPEUTIC TARGET

Jacob New[1], Levi Arnold[2], Megha Ananth[3], Sameer Alvi[3], Wen-Xing Ding[4], and Sufi Mary Thomas[1,2,3]

[1]Departments of Anatomy & Cell Biology; [2]Department of Cancer Biology; [3]Department of Otolaryngology; [4]Department of Pharmacology & Toxicology, University of Kansas Medical Center

Despite therapeutic advances, the survival rate for head and neck squamous cell carcinoma (HNSCC) has remained unchanged for the last fifty years. The HNSCC microenvironment includes various types of non-cancerous cells. Tumor associated fibroblasts (TAFs) are one such cell type that constitute a large portion of HNSCC tumors. Our lab and others have demonstrated the profound tumor promoting role of TAFs. TAFs exhibit more vesicular cytoplasm than normal fibroblasts from cancer-free patients (NFs). Further studies revealed these vesicles are most likely autophagosomes. Autophagosomes are formed

FEMTOSECOND LASER NANOTEXTURING OF DRUG-ELUTING STENTS

Mahmood Al Bashir and Rajeev Nair
Department of Mechanical Engineering, Wichita State University

Stents have been used for over a decade to treat coronary arteries affected by stenosis, a state in which the arteries are narrowed by the deposition of plaque. However, studies have shown that a few years after placement of a stent, restenosis can occur at the site, hindering its purpose. Drug-eluting stents (DES) have a significant impact in treating in-stent restenosis. DES are embedded with drugs like Sirolimus that are released over time once the stent is placed within the artery, reducing the chance of plaque deposition around the stent. But DES increases the risk of thrombosis, formation of blood clots around the stent due to a change in the blood flow profile. To improve the DES characteristics, femtosecond pulsed lasers are used to create nano-textures on the metallic stents. Nano-textures increase the availability of the drug, increase adhesion of the drug and the stent, and minimize requirement of a polymer carrier. If the nano-textured materials can show effective fluid retention characteristics, this can be a cost-effective solution to reduce or completely eliminate in-stent restenosis and thrombosis without the need of post-finishing operations. The interaction of bio-elements with the metallic stents can also be tested to ensure the efficiency of the DES.

REDUCING PTSD FOR VETERANS AND EMERGENCY SERVICE PROVIDERS

Daniel Clifford and Greg Meissen
Department of Psychology, Wichita State University

Post-traumatic stress disorder (PTSD) is a mental health diagnosis that can result from exposure to both stress and trauma. PTSD negatively impacts a disproportionate number of veterans and first responders and is related to a variety of negative health outcomes such as depression, anxiety, chronic health issues, and suicidality. These health issues cost the government billions of dollars each year. This project responded to this problem by helping design, implement, and evaluate a community-based, peer-led initiative to teach mind-body self-care skills to veterans and first responders in the Wichita, community. Veteran facilitators were trained and certified through the Center for Mind-Body Medicine to lead a 10-week intervention designed to reduce PTSD symptoms and increase overall functioning of participants. Interviews ete(d)-6 trenpa atok a-

“IF IT BLEEDS, IT LEADS” VS. #HESSTONSTRONG:
A FANTASY-THEME ct 8DC ne.a neQ0 0 eQe(y)-3 (YS4.8 (I)(M)-18 (E)3.9 3.0

DETERMINING AN EFFECTIVE TREATMENT PLAN FOR BREAST CANCER: A MULTI-CRITERIA DECISION MODEL AND ALGORITHM

Mostafa Hasan[1], [1], and Elshami Elamin[2]
[1]*Department of Industrial and Manufacturing Engineering, Wichita State University*
[2]*Central Care Cancer Center, Wichita, KS*

Breast cancer is the second leading cause of cancer deaths in U.S. women. For 2017, the American Cancer Society estimates that 255,180 women will be diagnosed with invasive breast cancer and 40,610 will die. Selection of an effective, patient-specific treatment plan has challenged physicians because the decision process involves many critical factors such as disease stage, disease risk factors, biomarker-related risks, and patient-related risks. In this study, a comprehensive set of criteria for selecting treatments was determined by literature review and oncologist interviews, and two analytical hierarchy process (AHP) models were used to weight criteria for both primary and secondary treatment therapies. Using the weighted criteria, we propose a treatment-ranking algorithm that evaluates every scenario and provides optimal patient-tailored treatment alternatives. We validate the multi-criteria ranking algorithm by comparing its treatment rankings

or is significantly correlated with the expert ranking. Thus the algorithm, on an easy-to-use Microsoft Excel platform, could be an accessible decision-support tool to aid oncologists and educate patients in choosing breast cancer treatment. Our multi-criteria ranking approach can also be adapted to solve complex decision-making problems in other service industries.

DIFFERENTIAL REGULATORY GENE EXPRESSION AT THE PROTEOMIC LEVEL IN SUBTYPES OF HUMAN OVARIAN CANCER CELL-LINES

Lipilekha Mukherjee, Isabel R. Hendry, and William J. Hendry
Department of Biological Sciences, Wichita State University

Ovarian carcinoma is the most lethal neoplasm and the fifth leading cause of mortality in women. Among the different subtypes, widely heterogeneous high-grade serous ovarian cancer (HGSOC) poses a great challenge to modern chemotherapy, due to its high recurrence rate and resistance to standard treatments. Recent studies suggest that survival time-line and disease recurrence are linked to the variable expression of biomarkers in cancer cells. Therefore, using Western blot analyses, we assessed the expression of a host of regulatory gene protein products at the whole-cell level in extracts from both three HGSOC and five non-HGSOC cell lines. Some of our most distinctive findings from those analyses were that: 1) Nrf2, a key factor in the anti-oxidant response element system, is expressed as a series of protein bands of variable molecular weight in all but the Kuramochi (HGSOC) cell line; 2) Estrogen receptor alpha is expressed in all but the OVSAHO (HGSOC) cell line; and 3) the AUF-1, p120-catenin, and NFKB-p50 proteins are expressed in both HGSOC and non-HGSOC cell types. We will next follow up those findings conducted at the whole-cell level with assessments at the individual cell and subcellular level using immunohistochemical analyses.

NUTRITION GUIDELINES IN KANSAS ELEMENTARY SCHOOLS

Erin Sawyers, Megan Spradlin, Jordan Heins, and Michelle Wallace, MPAS, PA-C
Department of Physician Assistant, Wichita State University

Childhood obesity is a nation-wide epidemic. Although the National School Lunch Program

ambiguous snack guidelines may be counteracting NSLP efforts. Our study assessed components of school lunches and snack guidelines in public elementary schools across Kansas. A question set was developed to evaluate these items. Answers were obtained from public information on school websites. Schools were qualitatively evaluated and compared based on size and socioeconomic status as defined in the study. Our data found that Kansas public elementary schools are following the NSLP guidelines; however, snack guidelines vary greatly. Our data do not identify any patterns in number of lunch meal options or snack guideline similarities amongst schools. A set of national or state snack guidelines may help to establish basic nutrition knowledge and encourage healthy eating in schools to combat the complex epidemic of childhood obesity.

CUBESAT PLATFORM FOR RESEARCH, EDUCATION AND BUSINESS IN KANSAS

Akshay Reddy Tummala, Suwat Sreesawet, and Atri Dutta
Department of Aerospace Engineering, Wichita State University

A CubeSat is a miniaturized satellite weighing 1

IMPROVING TRANSIT IN WICHITA: MARRYING THE OLD AND THE NEW

Mercy Umeri

Hugo Wall School of Public and Urban Affairs, Wichita State University

The transportation system in Wichita has served a diverse customer base over the years with its traditional system of large buses on fixed routes and smaller buses for specialized transportation (such as for the physically challenged, or elderly). Shrinking state revenues, increases in population and a more diverse set of needs have necessitated a push to find creative solutions to transit service delivery. Advances in technology have brought about variety and technology-enabled services that have increased choice in urban transit. Ride-sharing firms like Uber and Lyft employ a business model that combines personal services with technology. This business model fits the competitive

vetails seamlessly with the larger dynamics shaping

mix of services, the use of technology-based transportation in the United States and the transportation sector in Wichita, Kansas. A combination of the case study method, participant observation and empirical data will be used. Results from my studies will inform decision makers as they seek solutions to improve transit in the Wichita area.

NOTES:

Index of Presenters

Al Bashir, Mahmood 29
Bechtold, Rachel 19
Bhoyate, Sanket 19
Breeden, Joshua 26
Caudle, Keri 13
Christiano, Andrew 11
Clifford, Daniel 29
Coll, James 22
Emerson, Sam R. 14
Enninful, Regina 14
Fisher, Arianne 11
Flores, Lynzee 20
Frier, Ryan 8
Galliart, Matthew 15
Gardiepy, Kelsay E. 30
Hasan, Mostafa 31
Houk, Thomas 8
Huck ,Ian 27
Huffhines, Lindsay 22
Jennings, Jeffrey C. 23
Jorgensen, Darrah 12
Joshua Levi Weese 18
Karim, Md Zahidul 15
Keow, Alicia 30
Koch, H. Martin 23
Lemmon, Jack 16
Mukherjee, Lipilekha 31
Murphy, Maxwell D. 24
Nelson-Brantley, Heather V. 26
New, Jacob 27
Nguyen, Kien 24
Ralston, Mary M. 9
Ranaweera, C.K. 21
Rudeen, Aaron J. 25
Rusk, Adam 12
Sawyers, Erin 32
Schmid, Ryan 16
Shuguftha, Naz 20
Steele, Catherine 17
Thimmesch, Janelle 9
Tran, Tri V. 17
Tummala, Akshay Reddy 32
Umeri, Mercy 33
Weese, Joshua Levi 18
Wehrbein, Randol 10
Wisdom, Cate 25
Zarifa, Kelly 28
Zhao, Zheng 18